



Contents lists available at SciVerse ScienceDirect

Theoretical Computer Science

journal homepage: www.elsevier.com/locate/tcs

Kolmogorov complexity of initial segments of sequences and arithmetical definability

George Barmpalias*, C.S. Vlek

Institute for Logic, Language and Computation, Universiteit van Amsterdam, P.O. Box 94242, 1090 GE Amsterdam, The Netherlands

ARTICLE INFO

Article history:

Received 6 July 2010

Received in revised form 28 May 2011

Accepted 6 June 2011

G. Ausiello

Keywords:

Prefix-free Kolmogorov complexity

Relative randomness

 K -degrees

Arithmetical complexity

ABSTRACT

The structure of the K -degrees provides a way to classify sets of natural numbers or infinite binary sequences with respect to the level of randomness of their initial segments. In the K -degrees of infinite binary sequences, X is below Y if the prefix-free Kolmogorov complexity of the first n bits of X is less than the complexity of the first n bits of Y , for each n . Identifying infinite binary sequences with subsets of \mathbb{N} , we study the K -degrees of arithmetical sets and explore the interactions between arithmetical definability and prefix-free Kolmogorov complexity.

We show that in the K -degrees, for each $n > 1$, there exists a Σ_n^0 non-zero degree which does not bound any Δ_n^0 non-zero degree. An application of this result is that in the K -degrees there exists a Σ_2^0 degree which forms a minimal pair with all Σ_1^0 degrees. This extends the work of Csimá and Montalbán (2006) [8] and Merkle and Stephan (2007) [17]. Our main result is that, given any Δ_2^0 family \mathcal{C} of sequences, there is a Δ_2^0 sequence of non-trivial initial segment complexity which is not larger than the initial segment complexity of any non-trivial member of \mathcal{C} . This general theorem has the following surprising consequence. There is a $\mathbf{0}'$ -computable sequence of non-trivial initial segment complexity, which is not larger than the initial segment complexity of any non-trivial computably enumerable set.

Our analysis and results demonstrate that, examining the extend to which arithmetical definability interacts with the K reducibility (and in general any ‘weak reducibility’) is a fruitful way of studying the induced structure.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

The desire to compare the randomness ‘degree’ of two infinite binary sequences has led to the introduction of randomness reducibilities. An infinite sequence is called random if the prefix-free complexity K of its initial segments is very high, namely, at least as much as the length of the very segment (modulo a constant). Therefore a straightforward way to compare two sequences with respect to randomness is to compare the prefix-free complexity of their initial segments. Let $K(\sigma)$ denote the prefix-free complexity of string σ and say that $A \leq_K B$ if $K(A \upharpoonright_n) \leq^+ K(B \upharpoonright_n)$ for all $n \in \mathbb{N}$. By \leq^+ , we mean that the inequality holds modulo a constant that does not depend on n . This measure of randomness is called K -reducibility and was studied in [18], along with its plain Kolmogorov complexity counterpart. The induced structure of K -degrees has been a subject of interest in the past 5 years or so, though in terms of development this area is still in its infancy.

* Corresponding address: State Key Laboratory of Computer Science, Institute of Software, Chinese Academy of Sciences, Beijing 100190, P.O. Box 8718, People’s Republic of China.

E-mail addresses: barmpalias@gmail.com (G. Barmpalias), charlottevlek@gmail.com (C.S. Vlek).

URL: <http://www.barmpalias.net> (G. Barmpalias).

Table 1

Equivalence relations with respect to various weak reducibilities and their meanings (see [18] and [20, Section 8]).

\equiv_K	Same prefix-free complexity of the corresponding initial segments.
\equiv_C	Same plain complexity of the corresponding initial segments.
\equiv_{LK}	Same relativized prefix-free complexity.
\equiv_{LR}	Same notion of relativized randomness.

Miller and Yu studied the K -degrees of random sets in [18,19]. Csima and Montalbán constructed a minimal pair of K -degrees in [8]. Their method was highly non-constructive, making the pair merely Δ_4^0 (i.e. definable with four quantifiers), as noticed in [9, Section 10.13]. Merkle and Stephan (motivated by a number of related questions in [16]) studied the interaction between the Turing and the K reducibility in [17], along with its plain complexity counterpart, the C reducibility. Amongst many other results, they showed that there is a pair of Σ_2^0 sets which form a minimal pair in the K -degrees.

The study of the K reducibility is part of a larger study of the so-called ‘weak reducibilities’. These are preorders that measure various notions related to randomness (of sets), as opposed to computational complexity. Such reducibilities, like K , do not have an underlying map, i.e. an algorithm mapping (reducing) the second set to the first one. The existence of such maps is a vital feature in the Turing or stronger reducibilities.

In the Turing degrees, Post’s theorem gives an important link between reducibility (computability) and definability. For example, if a set is Turing reducible to a Σ_1^0 set then it is Δ_2^0 . A lot of methods that underlie the theory of Turing degrees rest on this link with definability. This breaks down when one considers weak reducibilities. For example, a feature that one finds in most weak reducibilities is that they can have uncountable lower cones. That is, there are uncountable classes, all of whose elements are reducible to a single set. Consider a related weak reducibility that was defined in [20], the LK reducibility (see Table 1). We say that $A \leq_{LK} B$ if $K^B(\sigma) \leq^+ K^A(\sigma)$ for all strings σ , where K^X denotes the prefix-free complexity relative to X , i.e. if B compresses more than (or at least as well as) A . It was shown in [6]¹ that for sufficiently ‘strong’ oracles B , the \leq_{LK} -cone below B is uncountable. Such properties also affect the study of local structures of the degrees, for example restricted to the Σ_1^0 or the Δ_2^0 sets. To illustrate this, consider a Δ_2^0 set B and $A \leq_T B$; then A is Δ_2^0 . However, by [2], there are uncountably many A such that $A \leq_{LK} B$ (unless $B \leq_{LK} \emptyset$). This means that there is no hope to derive any definability of A from B when $A \leq_{LK} B$. (Having said that, in the special case where the halting probability Ω is random relative to B , the relation $A \leq_{LK} B$ implies that A is Δ_2^0 relative to B . This was shown in [15].)

As a result, a number of methods that we use in the study of the Turing degrees do not have a counterpart in the study of ‘weaker’ degrees. Following up such differences sometimes lead to elementary differences between classical structures like the Turing degrees and related structures based on weaker reducibilities. For a number of such examples, we refer to [1].

But why is it useful to look for definability in weaker reducibilities? The presence of definability in a weak reducibility indicates that methods from the classical theory of the Turing degrees may be applicable for its study. We illustrate this by an analysis of definability in the K -degrees which, amongst other things, gives new ways to obtain minimal pairs in this structure. In Section 2, we study the class of *infinitely often* (i.o.) K -trivial sequences. These are sequences that have infinitely many initial segments σ with the property that $K(\sigma) \leq^+ K(|\sigma|)$ (throughout the paper we may identify $K(|\sigma|)$ with $K(0^{|\sigma|})$ so that we restrict ourselves to machines that output strings and not numbers). We note that the class of sets that are \leq_K -below such sequences is very well behaved; in particular, it is countable. Therefore such sequences locally generate good definability conditions. We also show that these sequences are rather common. Every truth table degree contains an i.o. K -trivial set; in particular, they are uncountably many. Also every (weakly) 1-generic set is i.o. K -trivial, so they form a co-meager class.

In Section 3, we study how arithmetical complexity interacts with the structure of K -degrees. Given a degree structure, the Σ_1^0 degrees are the ones which contain a Σ_1^0 set. The same applies to other classes of the arithmetical hierarchy. We show that, in the K -degrees, for each $n > 1$ there exists a non-zero Σ_n^0 degree which does not bound any non-zero Δ_n^0 degree. The particular case $n = 2$, combined with the basic properties of the i.o. K -trivial sets from Section 2, gives a Σ_2^0 degree which forms a minimal pair with every non-zero Σ_1^0 degree. This extends the work of Csima/Montalbán [8] and Merkle/Stephan [17] on minimal pairs in the K -degrees. However, their methods are entirely different from ours.

In fact, it is possible in the K -degrees to construct a Δ_2^0 non-zero degree which does not bound any Σ_1^0 non-zero degree. This result requires more effort and is rather surprising as Σ_1^0 sets have relatively low initial segment complexity. It also shows a contrast between the local structures of the K and the LK degrees, since in [1], it was shown that in the LK degrees every non-zero Δ_2^0 degree bounds a non-zero Σ_1^0 degree. Our method shows that, more generally, given any uniformly $\mathbf{0}'$ -computable family of sets there exists a $\mathbf{0}'$ -computable set of non-zero K -degree such that no set in the family is \leq_K -reducible to it, unless it is reducible to \emptyset . The proof of this main result is presented in Section 4.

The first construction of a minimal pair in the K -degrees was given in [8] through a brute-force argument. The proof relied on the construction of a non-decreasing unbounded function f such that for each set X there exists a constant c with

¹ In this paper, the result is proved for the related reducibility \leq_{LR} (see Table 1 for the meaning of the induced equivalence relation \equiv_{LR}) but by [13, Corollary 2.7] the reducibilities \leq_{LR} and \leq_{LK} coincide.

the property

$$\forall n [K(X \upharpoonright_n) \leq K(n) + f(n) + c] \iff X \leq_K \emptyset. \tag{1.1}$$

We refer to functions with the above property as *gap functions for K -triviality* and study them in Section 5. For example, we show that there is no Δ_2^0 unbounded non-decreasing gap function for K -triviality. This shows that the method used in [8] cannot be used in order to produce minimal pairs in the K -degrees of arithmetical complexity less than Σ_2^0 . Gap functions for K -triviality are interesting in their own right and are also related to the so-called Solovay functions that were studied in [4,11]. In Section 5, we study their arithmetical complexity and discuss the role they play in the K -degrees.

2. Infinitely often K -trivial sets

A set A is called *low for K* if the compression of strings is not improved when A is used as an oracle. In other words, if $K^A \equiv^+ K$. Here we say that $f \equiv^+ g$ for two functions f, g if $f \leq^+ g$ and $g \leq^+ f$. Hirschfeldt and Nies showed in [20] that lowness for K is equivalent to K -triviality. In [15], Miller defined a weak version of lowness for K by requiring that $K^A(n) \equiv^+ K(n)$ for infinitely many n (instead of all n). Such oracles A are known as *weakly low for K* . This variation turned out to be a fruitful characterization of another notion which is known as *lowness for Ω* . A set is called *low for Ω* if the latter is Martin-Löf random relative to it. Moreover, it turned out that within the class Δ_2^0 , an oracle is weakly low for K if and only if it is low for K . Consider the following analogous weakening of the notion of K -triviality.

Definition 2.1. A set A is called K -trivial on a set $M \subseteq \mathbb{N}$ with constant c if $K(A \upharpoonright_n) \leq K(n) + c$ for all $n \in M$. If it is K -trivial on an infinite set, then we call it *infinitely often K -trivial with constant c* .

A simple argument in [21, Exercise 5.2.9] shows that K -triviality on an infinite computable set coincides with K -triviality. In the following, we show that the class of infinitely often K -trivial sets is rather large, and quite different to the class of weakly low for K sets. Recall that given an enumeration of a set in stages, there are infinitely many n, s such that n is enumerated at stage s and no $i < n$ is enumerated at any stage $r \geq s$. Given a c.e. set A (and a computable enumeration of it with no repetitions), let us call the set of all such n (which are part of a pair n, s as above) a *set of minimal enumerations* of A . The following proposition was shown for plain complexity in [11] using the same argument. Moreover, it has been known to a number of researchers, although we are not aware of any explicit reference in the literature.

Proposition 2.2. *Every c.e. set is infinitely often K -trivial (on the set of its minimal enumerations).*

Proof. Fix a computable enumeration (A_s) of A without repetitions and a universal prefix-free machine U . Machine M does the following for each $n \in \mathbb{N}$. It waits for a stage s where n is enumerated in A and assigns to $A_s \upharpoonright_n$ all U -descriptions of 0^n . Since each number is enumerated in A at most once, M is prefix-free. If n is a minimal enumeration of A it is clear that $K_M(A \upharpoonright_n) \leq K(0^n)$. Hence $K(A \upharpoonright_n) \leq K(n) + c$ for some constant c and all n in the set of minimal enumerations of A . \square

The following results show that the sets that are \leq_K -below an infinitely often K -trivial set Y are Δ_2^0 definable in Y .

Proposition 2.3. *Suppose that Y is infinitely often K -trivial. Then each set in the lower cone $\{X \mid X \leq_K Y\}$ is computable in $Y \oplus \emptyset'$.*

Proof. Suppose that Y is infinitely often K -trivial via constant c_0 and $X \leq_K Y$ via c_1 . Let $c = c_0 + c_1$ and $F_c(n) := \{\sigma \mid |\sigma| = n \wedge K(\sigma) \leq K(n) + c\}$. By the coding theorem, we have that there is some constant b such that $|F_c(n)| < 2^{c+b}$ for all $n \in \mathbb{N}$. Since the prefix-free complexity function K is computable from \emptyset' , the infinite set M on which Y is K -trivial (via constant c_0) is computable from $Y \oplus \emptyset'$. Hence the downward closure of the set of strings $\cup_{n \in M} F_c(n)$ is computable from $Y \oplus \emptyset'$. Let us denote this tree by L_c . The cardinality of the levels of L_c have the same constant bound 2^{c+b} . By the choice of c , the set X is an infinite path through L_c . Since L_c is a $Y \oplus \emptyset'$ -computable tree with a constant bound on the cardinality of its levels, its infinite paths are computable in $Y \oplus \emptyset'$. \square

Proposition 2.4. *If Y is K -trivial on an infinite set M , then it is computable from $\emptyset' \oplus M$.*

Proof. The tree L_c from the proof of Proposition 2.3 is also computable in $\emptyset' \oplus M$. Since there is a constant bound on the cardinality of its levels, its paths (including Y) are computable in $\emptyset' \oplus M$. \square

By Proposition 2.3, every set that is \leq_K -below an infinitely often K -trivial Δ_2^0 set Y is Δ_2^0 . However, we do not know if the class of sets that are \leq_K -below Y is (uniformly) Δ_2^0 . To be more precise, we recall the following definition from computability theory. Let (Φ_e) be an effective list of all Turing functionals.

Definition 2.5. A class \mathcal{C} of subsets of \mathbb{N} is called a Δ_2^0 family (or uniformly \emptyset' -computable) if it can be written in the form $\{C_e \mid e \in \mathbb{N}\}$ where $C_e = \{n \mid \psi(e, n)\}$ and ψ is a Δ_2^0 property (i.e. a property that can be expressed in arithmetic with equivalent Σ_2^0 and Π_2^0 formulas). Equivalently, if there is a computable function f such that $\mathcal{C} = \{\Phi_{f(e)}^{\emptyset'} \mid e \in \mathbb{N}\}$, where $\Phi_{f(e)}^{\emptyset'}$ is total for each $e \in \mathbb{N}$.

Recall that a set is ω -c.e. if there is a computable function g and a computable approximation of it such that, for each $n \in \mathbb{N}$ the number of changes of the n th digit during the approximation is bounded by $g(n)$. It is not hard to see that the ω -c.e. sets form a Δ_2^0 family while the Δ_2^0 sets do not. A basic fact about the K -trivial c.e. sets is that they form a uniformly c.e. family

of sets (e.g. see [21, Fact 5.2.6]). Perhaps more interestingly, the K -trivial sets form a Δ_2^0 family. This follows from the fact that the ω -c.e. K -trivial sets form a Δ_2^0 family (see [21, Theorem 5.3.28]) and the deeper fact that K -trivial sets are ω -c.e. (see [21, Corollary 5.5.4]). In particular, the lower cone in the K -degrees below \emptyset is a Δ_2^0 family. We do not know if there are non-trivial lower cones in the K -degrees with the same property. The notions introduced in Definition 2.5 will play an important role in Sections 3 and 4.

In terms of Lebesgue measure the class of infinitely often K -trivial sets is small (i.e. it has measure 0). Indeed, no infinitely often K -trivial set is Martin-Löf random. However, in most other respects it is rather large, as we demonstrate below. We first need the following fact.

Lemma 2.6. *Let V be an infinite c.e. set with the property that for each $n \in \mathbb{N}$ there is at most one string of length n in V . Then $K(\sigma) \leq^+ K(|\sigma|)$ for all $\sigma \in V$.*

Proof. Let U be the universal prefix-free machine. Consider a prefix-free machine which, given $\sigma \in V$ it assigns to σ the U -descriptions of $0^{|\sigma|}$. By the properties of V such a machine exists, and $K(\sigma) \leq^+ K(0^{|\sigma|}) \leq^+ K(|\sigma|)$ for each $\sigma \in V$. \square

A tree T (as a downward closed set of binary strings) is called *pruned* if it has no dead-ends. In other words, if every $\sigma \in T$ has an extension in T .

Theorem 2.7. *There is a computable pruned perfect tree such that every path in it is infinitely often K -trivial. In particular, there are 2^{\aleph_0} infinitely often K -trivial sets.*

Proof. Consider a computable tree $T : 2^{<\omega} \rightarrow 2^{<\omega}$ such that $|T(\sigma)| \neq |T(\tau)|$ if $\sigma \neq \tau$. Then the range of T is a c.e. set V which satisfies the properties of Lemma 2.6. Hence any path $\cup_n T(X \upharpoonright_n)$ through the tree is K -trivial on the infinite set of numbers $|T(X \upharpoonright_n)|$, $n \in \mathbb{N}$. \square

The following Corollary is implicit in [17], although it is obtained using different methods.

Corollary 2.8. *Every truth table degree contains an infinitely often K -trivial set.*

Proof. Let A be any set and let T be the tree of Theorem 2.7. Note that T can be viewed as a total Turing functional $\Phi : 2^\omega \rightarrow 2^\omega$ via $X \rightarrow \cup_n T(X \upharpoonright_n)$. Moreover, we can define a total Turing functional $\Psi : 2^\omega \rightarrow 2^\omega$ which is the inverse of Φ on the paths of T and some finite set on other paths. Now let $\Phi^A = B$, so that $\Psi^B = A$. Then $B \leq_{tt} A$ and $A \leq_{tt} B$. Moreover B is infinitely often K -trivial since it is on T . \square

Theorem 2.9. *Every (weakly) 1-generic set is infinitely often K -trivial. In particular, the class of infinitely often K -trivial sets is co-meager.*

Proof. Every weakly 1-generic set meets every infinite c.e. dense set of strings infinitely often. This is because every co-finite subset of a dense set of strings is dense. Hence, by Lemma 2.6 it suffices to define a dense set V of strings σ such that no two strings in V have the same length. Indeed, in that case we have $K(\sigma) \leq^+ K(|\sigma|)$ for each string $\sigma \in V$. Hence any sequence that intersects V is infinitely often K -trivial.

The set V is defined recursively as follows, based on a computable enumeration (σ_s) of all strings. We order the set of binary strings first by length and then lexicographically. At stage $s + 1$, put in V the least string which extends σ_s and its length is larger than the lengths of all the strings in $V[s]$. Clearly, V has the desired properties. \square

The following theorem can be combined with various basis theorems for Π_1^0 classes to give infinitely often K -trivial sets with additional properties.

Theorem 2.10. *There is a nonempty Π_1^0 class which consists of infinitely often K -trivial sets but does not contain any K -trivial sets.*

Proof. We only give a sketch of the proof since it does not involve new ideas. Let V be a c.e. dense set of strings σ such that $K(\sigma) \leq^+ K(|\sigma|)$. This is obtained as in the proof of Theorem 2.9. Let c be a constant such that $K(n) \leq 2 \log n + c$ for all n , where $\log n$ is the largest number k such that $2^k \leq n$. To avoid the K -trivial sets in the class we use a computable function f such that for all n and all strings σ of length n , there is an extension τ of σ of length $f(n)$ such that $K(\tau \upharpoonright_k) > 2 \log k + c + n$ for some $k < |\tau|$.

At step 1, we put all strings of length $f(1)$ in our tree and promise to remove any such σ which satisfies $K(\sigma \upharpoonright_k) \leq 2 \log k + c + 1$ for all $k \leq |\sigma|$ (this is a Π_1^0 event). Note that by this action, we also remove those σ such that $K(\sigma \upharpoonright k) \leq K(k) + 1$ for all $k \leq |\sigma|$. Let $\ell_1 = f(1)$. At step 2, for each of the chosen strings of step 1, we choose an extension τ in V . If ℓ is the length of the largest such extension, we let $\ell_2 = \ell$. We put on the tree each such extension τ concatenated with $\ell - |\tau|$ zeros. We also declare any other extension of σ that is incompatible with τ not to be part of the tree. We continue in the same way for the rest of the steps, where at step $2n + 1$ we put on T the extensions of the strings of step $2n$ of length $f(\ell_{2n})$. Also, we promise to remove those strings such that $K(\sigma \upharpoonright_k) \leq 2 \log k + c + n$ for all $k \leq |\sigma|$.

This procedure defines a Π_1^0 tree T such that the set $[T]$ of its paths is nonempty. Any real in $[T]$ intersects V infinitely often by the construction of T (in particular the even steps). Hence, it is infinitely often K -trivial. Moreover, it does not contain K -trivial reals by the odd steps of the construction. \square

If we combine [Theorem 2.10](#) with the computably dominated basis theorem (e.g. see [[21](#), Theorem 1.8.42]), we get that there are non-computable i.o. K -trivial computably dominated sets. This contrasts the fact that every computably dominated K -trivial set is computable. We close this section with two more subclasses of the infinitely often K -trivial sets. Recall that f is a DNC (or diagonally non-computable) function if $f(e) \neq \varphi_e(e)$ for all e such that $\varphi_e(e) \downarrow$ (where (φ_e) is an effective enumeration of all partially computable functions).

Theorem 2.11. *If there is no DNC function f such that $f \leq_{tt} A$ then A is i.o. K -trivial.*

Proof. Suppose that every function that is truth table reducible to A fails to be diagonally non-computable. Consider the function f which, given n , it outputs a code $\langle A \upharpoonright_n \rangle$ of the first n bits of the characteristic function of A . Since $f \leq_{tt} A$, f is not DNC. Therefore, $\varphi_e(e) \downarrow = \langle A \upharpoonright_e \rangle$ for infinitely many $e \in \mathbb{N}$. But for some constant c and each e such that $\varphi_e(e) \downarrow$, we have $K(\varphi_e(e)) \leq K(e) + c$. Since $K(\langle A \upharpoonright_e \rangle) \leq K(\langle A \upharpoonright_e \rangle) + b$ for some constant b and all $e \in \mathbb{N}$, there are infinitely many e such that $K(\langle A \upharpoonright_e \rangle) \leq K(\langle A \upharpoonright_e \rangle) + c + b$. Hence A is i.o. K -trivial. \square

According to [[12,14](#)], we say that a set A is *complex* if there is an unbounded non-decreasing computable function f such that $K(A \upharpoonright_n) \geq f(n)$ for all $n \in \mathbb{N}$. In the same paper, the authors showed that a set A is complex iff there is a DNC function f such that $f \leq_{wt} A$. Hence the following is a consequence of [Theorem 2.11](#).

Corollary 2.12. *If a set is not infinitely often K -trivial, then it is complex.*

Note that the converse of [Corollary 2.12](#) does not hold, since \emptyset' is complex but also i.o. K -trivial by [Proposition 2.2](#).

Recall that by [[10](#)] (also see [[9](#), Theorem 1.23.18]), if a set that is computed by a 1-generic then it does not compute a DNC function. Therefore, [Theorem 2.11](#) implies the following.

Corollary 2.13. *Every set that is computed by a 1-generic is i.o. K -trivial.*

[Theorem 2.11](#) shows that there are non-trivial lower cones in the Turing degrees that consist entirely of i.o. K -trivial sets. However i.o. K -trivial sets are not closed downward under Turing reducibility. Indeed, the halting set is i.o. K -trivial but it computes random sets.

3. Arithmetical complexity in the K -degrees

In this section, we explore the definability restrictions in \leq_K -lower cones. A consequence of this analysis is that there is a Σ_2^0 set which forms a minimal pair with any (non-trivial) c.e. set in the K -degrees. We start with the following, which has an analog in the Turing degrees. Moreover, the proofs in the two cases are similar.

Theorem 3.1. *There exists a Σ_2^0 set $A >_K \emptyset$ such that $X \not\leq_K A$ for all Δ_2^0 sets $X >_K \emptyset$.*

Proof. We will enumerate A in a \emptyset' -computable construction, so that A is $\Sigma_1^0(\emptyset')$, hence Σ_2^0 . To ensure that $A \not\leq_K \emptyset$, we need to meet the following requirements:

$$R_e : \exists n [K(A \upharpoonright_n) > K(n) + e].$$

To ensure that A does not K -bound any non-trivial Δ_2^0 sets, we meet the following:

$$N_e : [\Phi_e^{\emptyset'} \text{ is total and } \Phi_e^{\emptyset'} \not\leq_K \emptyset] \Rightarrow \exists n [K(\Phi_e^{\emptyset'} \upharpoonright_n) \not\leq K(A \upharpoonright_n) + e].$$

Note that if $K(\Phi_k^{\emptyset'} \upharpoonright_n) \leq K(A \upharpoonright_n) + t$ for all n and some $k, t \in \mathbb{N}$, then there is some $e > t$ such that $\Phi_e = \Phi_t$. Hence $K(\Phi_e^{\emptyset'} \upharpoonright_n) \leq K(A \upharpoonright_n) + e$ for all n . So requirements N_e are sufficient.

Suppose that we only wish to satisfy a single N_e (and all R_i). We can compute a constant c such that $K(1^n) \leq K(n) + c$ for all $n \in \mathbb{N}$. Fix a Martin-Löf random sequence $Y \leq_T \emptyset'$. We can proceed by defining $A \upharpoonright_s [s] = Y \upharpoonright_s$ while constantly searching for some n such that $\Phi_e^{\emptyset'} \upharpoonright_n \downarrow$ and $K(\Phi_e^{\emptyset'} \upharpoonright_n) > K(n) + c + e$. If we find such a number n at stage s , we enumerate all numbers $\leq n$ into A thus meeting N_e . In this case, for the positions $> n$ of A , we copy the corresponding digits of Y . If the search does not halt during the stages s , N_e is satisfied trivially. In any case, all R_i are met as A will be equal to Y apart from finitely many positions.

We combine these strategies for N_e , $e \in \mathbb{N}$ in order to construct A which satisfies all of these requirements. Each strategy N_e imposes a restraint $r_e[s]$ on A at stage s . The restraint imposed by N_e will restrict the lower priority strategies N_i , $i > e$ from changing certain (finite) segments of A . This will mainly help the satisfaction of the R_e requirements. On the other hand, the restraints will reach a limit, so that each N_e can work without interference from the higher priority requirements N_j , $j < e$ from some stage on. Let $c_e[s]$ be a constant such that $K(Z_e[s] \upharpoonright_n) \leq K(n) + c_e[s]$ for all $n \in \mathbb{N}$, where $Z_e[s] = (A \upharpoonright_{r_{e-1}})[s] * 1^\omega$. Here $*$ denotes concatenation and $r_{-1}[s] := 0$ for all s . Note that $c_e[s]$ is computable from $(A \upharpoonright_{r_e})[s]$. Set $r_e[0] = 0$ for all e . If $r_e[s + 1]$ is not defined explicitly in the construction, we have $r_e[s + 1] = r_e[s]$. We say that N_e *requires attention at stage* $s + 1$ if there exists $n \leq s$ such that $\Phi_e^{\emptyset'} \upharpoonright_n \downarrow$ and $K(\Phi_e^{\emptyset'} \upharpoonright_n) > K(n) + c_e[s] + e$.

Construction. At stage $s + 1$, let $m_s = \max\{s, \max A[s]\}$, where $\max A[s]$ denotes the largest element in the finite set $A[s]$. Also, find the least $e < s$ such that N_e requires attention and is not currently declared *satisfied*. Enumerate all numbers n

with $r_{e-1}[s] < n \leq m_s$ into A . Define

$$A[s + 1] = (A \upharpoonright_{r_{e-1}})[s] * 1^{m_s - r_{e-1}[s]} * Y \upharpoonright_k$$

where k is the least number such that $K(A[s + 1] \upharpoonright_{m_s+k}) > K(m_s + k) + e$. Finally set $r_e[s + 1] = m_s + k$, declare N_e satisfied and all $N_i, i > e$ not satisfied. If no $N_e, e < s$ requires attention, let $A[s + 1] = A[s] * Y(s)$, where $Y(s)$ is the s th digit of Y .

Verification. If only finitely many N_e require attention during the construction, $r_i, c_i, i \in \mathbb{N}$ reach a limit and $A = \sigma * Y_*$ for some string σ and a final segment Y_* of Y . Hence all R_e are satisfied. Moreover, if some N_e was not satisfied, it would require attention at some stage of the construction. Hence almost all (therefore, by the padding lemma, all) N_e are satisfied.

If infinitely many N_e require attention during the construction, infinitely many of them receive attention. If at some stage $s + 1$ requirement N_e receives attention and no $N_i, i < e$ receives attention after $s + 1$, requirement R_e is satisfied (and remains so for the rest of the stages). Indeed, by the choice of k and the definition of r_e in the construction, we have $K(A[s + 1] \upharpoonright_{r_e}) > K(r_e) + e$, the restraint r_e has reached a limit at stage $s + 1$ and $A[s + 1] \upharpoonright_{r_e} = A \upharpoonright_{r_e}$. Since infinitely many N_e require attention during the construction, there will be infinitely many stages $s + 1$ when some N_e receives attention and no $N_i, i < e$ ever requires attention after stage $s + 1$. Therefore by the above observation infinitely many R_e are met. This in turn implies that all R_e are met.

Finally, we show by induction on e that each N_e is satisfied and r_e, c_e reach a limit. Suppose that this holds for $e < k$ and let s_0 be a stage after which the values of r_e, c_e remain constant for all $e < k$. If N_k does not require attention after stage s_0 , it is satisfied and r_e, c_e remain constant after s_0 . Otherwise N_k will receive attention at some stage $s_1 > s_0$ and will be satisfied according to the action taken in the construction (the definition of $(A \upharpoonright_{r_k})[s_1]$) and the fact that $A \upharpoonright_{r_k}$ will be preserved from then on. Note that N_k will not receive attention after stage s_1 , thus r_k, c_k reach a limit at that stage. This concludes the induction step and the proof. \square

The following result improves the complexity of the minimal pair of K -degrees that was constructed in [17].

Corollary 3.2. *There is a Σ_2^0 set whose greatest lower bound with every Σ_1^0 set is $\mathbf{0}$ in the K -degrees.*

Proof. Let A be the Σ_2^0 set of Theorem 3.1 and B any c.e. set. By Propositions 2.2 and 2.3, every set $X \leq_K B$ is Δ_2^0 . Therefore, if $X \leq_K A$ by the choice of A the set X has to be K -trivial. \square

Note that the argument we gave in the proof of Theorem 3.1 relativizes to $\emptyset^{(n)}$ for all $n > 0$, giving analogs on each level of arithmetical complexity. Hence we have the following.

Theorem 3.3. *Let $n > 1$. There exists a Σ_n^0 set $A >_K \emptyset$ such that $X \not\leq_K A$ for all Δ_n^0 sets $X >_K \emptyset$.*

As above, this gives the following application to the study of minimal pairs in the K -degrees.

Corollary 3.4. *Let $n > 1$. There exists a Σ_n^0 set $A >_K \emptyset$ whose greatest lower bound in the K -degrees with any Δ_n^0 infinitely often K -trivial set is $\mathbf{0}$.*

Proof. By Propositions 2.2 and 2.3, the lower cone below an i.o. K -trivial Δ_n^0 set consists entirely of Δ_n^0 sets. Hence the Σ_n^0 set of Theorem 3.3 has the desired properties. \square

Theorem 3.3 can be seen as a strong separation of the Σ_n^0 classes from their predecessors Δ_n^0 in the K -degrees. An immediate question is whether we can also separate Δ_n^0 from Σ_{n-1}^0 in the same way. In Section 4, we show the following.

Theorem 3.5. *Given any Δ_2^0 family of sets there exists a Δ_2^0 set whose K -degree is non-zero and does not bound any non-zero K -degree of a set in the family.*

Since the class of Σ_1^0 sets is a Δ_2^0 family, we get the following.

Corollary 3.6. *In the K -degrees, there is a Δ_2^0 non-zero degree that does not bound any Σ_1^0 non-zero degree.*

The above result is rather surprising as Σ_1^0 sets have relatively low initial segment complexity.

4. Proof of Theorem 3.5

Suppose that (X_e) is a uniformly \emptyset' -computable family of sets. To simplify the requirements, assume without loss of generality that each set in the family has infinitely many indices in this list. Let $X_e[s]$ be a computable system of approximations to the sets in the family. Then $K(X_e \upharpoonright_n)[s]$ (where the ‘suffix’ applies to both X_e and the complexity K , so that they are simultaneously computably approximated) is a computable system of approximations to their initial segment complexities. For each e we will make sure that the following requirements are satisfied:

$$R_e : \exists n [K(X_e \upharpoonright_n) \not\leq K(A \upharpoonright_n) + e] \vee \forall k [K(X_e \upharpoonright_k) \leq^+ K(k)].$$

To test the K -triviality of X_e , the construction will enumerate a c.e. set of strings V and use Lemma 2.6. We will make sure that for each n there is at most one string in V of length n . By Lemma 2.6, the satisfaction of R_e follows from the satisfaction

of the following modified requirement.

$$N_e : \begin{cases} \text{There exists a c.e. set } V \text{ as in Lemma 2.6 such that either for some } n \text{ we have} \\ K(X_e \upharpoonright_n) \not\leq K(A \upharpoonright_n) + e, \text{ or for all } \sigma \in V \text{ we have } K(X_e \upharpoonright_{|\sigma|}) \leq K(\sigma) + e. \end{cases}$$

Indeed, if the second clause of N_e holds, by Lemma 2.6 the set X_e will be K -trivial on an infinite computable set of lengths. Hence it will be K -trivial.

In the next section, we give an atomic construction which, given e , uniformly produces $A \leq_T \emptyset'$ which is not K -trivial and, if $X_e \not\leq_K \emptyset$ then $X_e \not\leq_K A$. Although this is not used explicitly in the main construction of Section 4.2, it helps understanding the ideas involved.

4.1. Strategy for one N_e

To increase the complexity of A we use a Δ_2^0 random set Y with computable approximation Y_s . Define $V = \{Y_s \upharpoonright_s \mid s \in \mathbb{N}\}$.

Let σ_s be the shortest string $\sigma \in V$ such that $|\sigma_s| \leq s$ and $K(X_e \upharpoonright_{|\sigma_s|})[s] > K(\sigma)[s] + e$. If this does not exist, let $\sigma_s = Y_s \upharpoonright_s$. The witness of the strategy at stage s is defined to be the string $\sigma_s * (Y_s \upharpoonright_s)$. Below, we show that the witnesses of the strategy in the various stages s converge to a unique infinite binary sequence. We define A to be this very sequence.

The set A converges. One of the following must occur.

- (a) The string σ_s reaches a (finite) limit τ .
- (b) The length of σ_s tends to infinity.

Indeed, if (a) does not hold we have that $K(X_e \upharpoonright_{|\sigma|}) \leq K(\sigma) + e$ for all $\sigma \in V$. In this case, each $\sigma \in V$ can only be chosen as σ_s finitely often. Therefore (b) must occur.

In the first case, there exists some stage s_0 such that the witness of the strategy is $\tau * (Y_s \upharpoonright_s)$ for all $s > s_0$. In this case, A converges to $\tau * Y$. In the second case, the witnesses converge to Y . Therefore A is well defined in any case.

The set A satisfies N_e and is not K -trivial. Clearly A is defined uniformly from the index e , a Δ_2^0 index of X_e , and \emptyset' . As explained above, in any case, Y is a tail of A . Therefore, A is not K -trivial. Finally, in case (a), we have $K(X_e \upharpoonright_{|\tau|}) > K(A \upharpoonright_{|\tau|}) + e$, since $\tau \subset A$. In case (b), we have that $K(X_e \upharpoonright_{|\sigma|}) \leq K(\sigma) + e$ for all $\sigma \in V$. By Lemma 2.6, we have that there is a constant c such that $K(\sigma) \leq K(|\sigma|) + c$ for all $\sigma \in V$. Hence X_e is K -trivial. Therefore, in any case the sets A, X_e satisfy N_e .

4.2. Satisfying all N_e

We will use a priority tree (the full binary tree) in order to construct A which meets all requirements. To make sure that A is not K -trivial, we need to meet the following requirements.

$$P_e : \exists n [K(A \upharpoonright_n) > K(n) + e].$$

Strategies are identified with nodes on the tree. Each node on the tree is 2-branching with outcomes $0 < 1$. For a node that is associated with N_e , the outcome 0 corresponds to the belief that X_e is K -trivial while outcome 1 corresponds to the negation of this belief. Along with the (current) outcome, each node will have a primary and a secondary witness. The primary witness will be as in Section 4.1, associated with the satisfaction of N_e . The secondary witness will be an extension of the primary witness that is associated with the satisfaction of P_e . The secondary witnesses will play the role that Y played in Section 4.1, i.e., they will increase the initial segment complexity of the constructed set A . In the following, whenever we refer to ‘witnesses’ of a strategy, we always mean both the primary and the secondary witness of it. Consider a computable partition of \mathbb{N} into infinite sets $\mathbb{N}^{[\alpha]}$ indexed by the strategies α . A node α will enumerate a c.e. set V_α containing strings of length in $\mathbb{N}^{[\alpha]}$. A strategy of length e on the leftmost infinitely often visited path (also called the true path) will run successfully and satisfy N_e, P_e .

In the following, we define the outcomes and witnesses of the strategies during the stages of the construction. Fix a \emptyset' -computable function (in both arguments) $p_e(\sigma)$, which gives some $\tau \supset \sigma$ such that $K(\tau) > K(|\tau|) + e$. Also let $p_e(\sigma)[s]$ be a computable approximation to it.

At stage s , a path δ_s of length s through the tree will be defined inductively, determining the ‘visited nodes’ at stage s . A β -stage is a stage s where β was visited, i.e. $\beta \subseteq \delta_s$. The root is the first visited node at each stage and the other visited nodes are determined by the current outcomes and witnesses of their predecessors. The outcome of a visited node α at stage s is 0 if $K(X_e \upharpoonright_{|\sigma|})[s] \leq K(\sigma)[s] + e$ for all strings $\sigma \in V_\alpha[s - 1]$ which extend the current witnesses of each $\beta \subset \alpha$. In this case, the primary witness of α is equal to the union of the current witnesses of each $\beta \subset \alpha$. Otherwise, the outcome is 1 and the primary witness of α is the shortest string $\sigma \in V_\alpha[s - 1]$, which extends the current witnesses of all $\beta \subset \alpha$ and $K(X_e \upharpoonright_{|\sigma|})[s] > K(\sigma)[s] + e$. In any case, the secondary witness of α is defined to be $p_e(\sigma)[s]$, where σ is its primary witness. Finally, the parameters of a node α are only updated at the α -stages.

4.3. Construction

At stage s , calculate the path δ_s of length s , starting from the root and following the current outcomes of the nodes. Pick the least number $n < s$, which is in some $\mathbb{N}^{[\alpha]}$ for $\alpha * 0 \subset \delta_s$ and such that there is no string of length n in V_α . Enumerate

into V_α the least string of length n , which is compatible with the current witnesses of δ_s . If such an n does not exist, go to the next stage.

4.4. Verification

Since the branching of the priority tree is finite, there exists a leftmost infinitely often visited infinite path δ . Moreover, it follows from the definition of the outcomes and witnesses that at each stage s , the witnesses of the initial segments of δ_s are linearly ordered.

Lemma 4.1. *Suppose that $\beta \subset \delta$ and α is the immediate predecessor of β . The witnesses of α reach a limit in the β -stages.*

Proof. The secondary witnesses are just the images of the primary witnesses under the Δ_2^0 function p . Therefore, it suffices to show the lemma for primary witnesses. We do this by induction on the length of α . Suppose that it holds for all $\alpha \subset \delta$ of length $< n$ and σ is the union of the final values of the witnesses of these nodes in the $\delta \upharpoonright_n$ -stages. We show that it holds for $\alpha = \delta \upharpoonright_n$. Let $\beta = \delta \upharpoonright_{n+1}$. If $\alpha * 0 \subset \delta$ the primary witness of α in the β -stages has limit σ . Otherwise $\alpha * 1 \subset \delta$, which means that the primary witness of α will reach a limit τ (over all stages) such that $K(X_e \upharpoonright_{|\tau|}) > K(\tau) + e$ for some $\tau \in V_\alpha$ and $e = |\alpha|$. \square

Given $\alpha \subset \delta$, the *true witnesses* of the immediate predecessor of α are defined to be the limits of its witnesses in the α -stages. In Lemma 4.3, we will define A to be the union of these true witnesses. Moreover, the *true outcomes* of the nodes on δ are the outcomes that lie on δ .

Lemma 4.2. *Suppose that $|\alpha| = e$. If $\alpha * 1 \subset \delta$ then V_α is finite and $K(X_e \upharpoonright_{|\sigma|}) > K(\sigma) + e$, where σ is the final witness of α . If $\alpha * 0 \subset \delta$ then V_α contains a string of each length in $\mathbb{N}^{|\alpha|}$ and X_e is K -trivial.*

Proof. For the first clause, note that if $\alpha * 1 \subset \delta$, then the construction will stop enumerating into V_α after some stage. Therefore, V_α is finite. Moreover, after some stage the primary witness of α will settle on the shortest string σ in V_α , which extends the true witnesses of its predecessors and $K(X_e \upharpoonright_{|\sigma|}) > K(\sigma) + e$.

For the second clause, suppose that $\alpha * 0 \subset \delta$. By the construction, the set V_α contains a string of each length in $\mathbb{N}^{|\alpha|}$. By Lemma 4.1, the witnesses of the predecessors of α reach a limit in the α -stages. Let σ be the union of these witnesses. By construction, almost all strings in V_α will be extensions of σ . Hence, the fact that $\alpha * 0 \subset \delta$ implies that for almost all $\tau \in V_\alpha$ (in particular, all that extend σ), we have $K(X_e \upharpoonright_{|\tau|}) \leq K(\tau) + e$. By Lemma 2.6, we have $K(\tau) \leq^+ K(|\tau|)$ for all $\tau \in V_\alpha$. Hence $K(X_e \upharpoonright_k) \leq^+ K(k)$ for almost all $k \in \mathbb{N}^{|\alpha|}$ and X_e is K -trivial. \square

The following lemma is crucial in that it enables us to define the set A and more importantly to ensure that it is Δ_2^0 .

Lemma 4.3. *The strings enumerated into the sets V_α during the construction converge to a unique sequence A , which is the union of the true witnesses of the nodes on δ . In other words, for all $n \in \mathbb{N}$ there exists a stage s_0 such that all strings enumerated by the construction after stage s_0 are extensions of $A \upharpoonright_n$.*

Proof. Let $\beta \subset \delta$ be a node with true secondary witness σ which reaches a limit in the $\delta \upharpoonright_{|\beta|+1}$ stages at stage s_* . In the following, all stages are assumed to be larger than s_* and the last stage where a node to the left of β was visited. Since β is an arbitrary initial segment of δ , the lemma is a consequence of the following.

Claim: There is a stage s_0 after which the only strings enumerated in the sets V_α (for all nodes α in the tree) are extensions of σ . (4.1)

Claim (4.1) clearly holds for the nodes α that lie on the left of β . Indeed, in this case, V_α is finite. By Lemma 4.1, it also holds for the nodes α that extend β and its true outcome. Indeed, in this case the strings enumerated in V_α must extend the current secondary witness of β , which reaches limit σ in the α stages. Finally it holds for the nodes $\alpha \subset \beta$ such that $\alpha * 1 \subset \delta$ since in this case, by Lemma 4.2, V_α is finite. Hence, it remains to show Claim (4.1) for the case where $\alpha \subseteq \beta$ and $\alpha * 0 \subset \delta$, or α is to the right of the true outcome of β . The latter case holds when α extends some $\eta * 1$ where $\eta \subseteq \beta$ and $\eta * 0 \subset \delta$.

In the latter case, the choice of these η implies that the length of their witnesses at stages s where $\delta_s \supset \eta * 1$ tends to infinity. So, if we show that almost all strings of V_η extend σ , we have that at almost all stages s such that $\delta_s \supset \eta * 1$ the witnesses of η extend σ . From this it follows that beyond some stage, any string enumerated to some V_α for $\alpha \supset \eta * 1$ must extend σ .

Hence, it remains to show Claim (4.1) for the particular case where $\alpha \subseteq \beta$ and $\alpha * 0 \subset \delta$. We prove this by finite induction. Let $\eta_0 \supset \dots \supset \eta_t$ be the descending sequence of all strings $\eta \subseteq \beta$ such that $\eta * 0 \subset \delta$. Fix $i < t$, suppose that the claim holds for all $\eta_j, j < i$ and let ρ_j be the union of the true witnesses of the predecessors of η_j (for each $j < i$). Also, let s_i be a stage beyond which we have $K(X_{e_j} \upharpoonright_{|\tau|}) \leq K(\tau) + e_j$ (where $e_j = |\eta_j|$) for each $j < i$ and each string τ in V_{η_j} , which is an extension of ρ_j but not an extension of σ . By induction hypothesis, there are finitely many such strings τ , so s_i exists.

If a string is enumerated in V_{η_i} at a stage $s > s_i$, then either δ_s extends the true outcome of β or $\delta_s \supset \eta_j * 1$ for some $j < i$. In the first case, the enumerated string must be an extension of the witness σ of β . In the second case, it must extend the current witness of some $\eta_j, j < i$ where $\delta_s \supset \eta_j * 1$. According to the choice of s_j , the current witness of η_j at stage s must extend σ . Hence, in either case the enumerated string is an extension of σ . This concludes the induction, the proof of Claim (4.1) and the proof of the lemma. \square

Lemma 4.4. *The set A is Δ_2^0 and satisfies all N_e, P_e for $e \in \mathbb{N}$.*

Proof. Lemma 4.3 shows how to calculate A by asking Σ_1^0 questions. Hence A is Δ_2^0 . Let $e \in \mathbb{N}$ and let α be the unique node on δ of length e . Also, let σ be the true primary witness of α .

For P_e it suffices to show that there is some string $\tau \subset A$ such that $p_e(\tau) \subset A$. Clearly $\sigma \subset A$ and $p_e(\sigma)$ is the secondary witness of α . Hence $p_e(\sigma) \subset A$ and P_e is satisfied. For N_e , suppose that X_e is not K -trivial. By Lemma 4.2, we have that $\alpha * 1 \subset \delta$ and $K(X_e \upharpoonright_{|\sigma|}) > K(\sigma) + e$. By the definition of A , we have $\sigma \subset A$ so N_e is satisfied. \square

An interesting feature of the above proof of Theorem 3.5 is that, although determining completely the outcomes of the strategies requires \emptyset'' (it is an infinite injury argument after all) the set A that we produce is computable in \emptyset' . In fact, we paid special effort to ensure the computable approximability of A in Lemma 4.3. This interesting feature is already apparent in the atomic module that we presented in Section 4.1. It would be desirable to replace this combinatorial argument with a \emptyset' -oracle construction determining A explicitly (i.e. not as the limit of a computable approximation).

5. Gap functions for K -triviality

An interesting fact from [8] is the existence of a non-decreasing unbounded function that can replace the constant in the definition of K -triviality. In this section, we isolate this notion and exhibit its role in the structure of the K -degrees. It is instructive to compare the results of this section with [19, Sections 3, 5], where a different notion of a ‘gap function’ plays a crucial role in analyzing the downward and upward oscillations of the initial segment prefix-free complexity of random sets.

Definition 5.1. We say that $f : \mathbb{N} \rightarrow \mathbb{N}$ is a *gap function for K -triviality* if for each set X , we have

$$\exists c \forall n [K(X \upharpoonright_n) \leq K(n) + f(n) + c] \iff X \text{ is } K\text{-trivial.} \tag{5.1}$$

Moreover, f is a *gap function for K -triviality of Δ_2^0 sets* if (5.1) holds for all Δ_2^0 sets X . An analogous definition holds for the other arithmetical classes.

If $\forall n [K(X \upharpoonright_n) \leq K(n) + f(n) + c]$, we say that X *obeys f* with constant c . Clearly the ‘ \iff ’ of the equivalence in Definition 5.1 holds always. An analysis of the construction of the gap function for K -triviality by Csima and Montalbán [8] shows the following.

$$\text{There is a } \Delta_4^0 \text{ unbounded and non-decreasing gap function for } K\text{-triviality.} \tag{5.2}$$

A simple analysis of the main argument in [8] shows the following connection between gap functions of K -triviality and minimal pairs in the K -degrees.

$$\begin{aligned} &\text{Let } f \text{ be any unbounded and non-decreasing gap function for } K\text{-triviality.} \\ &\text{Then } f \oplus \emptyset' \text{ computes two sets that form a minimal pair in the } K\text{-degrees.} \end{aligned} \tag{5.3}$$

Fact (5.3) also shows why the particular case of unbounded and non-decreasing gap functions is of special interest. The following converse of (5.3) also holds.

$$\begin{aligned} &\text{If } X, Y \text{ form a minimal pair in the } K\text{-degrees, then} \\ &f(n) := \min\{K(X \upharpoonright_n), K(Y \upharpoonright_n)\} - K(n) \\ &\text{is a gap function for } K\text{-triviality.} \end{aligned} \tag{5.4}$$

The following fact is useful in Theorem 5.2.

$$\text{If } f \text{ is a } \Delta_2^0 \text{ non-decreasing unbounded function, then there is an unbounded non-decreasing function } g \text{ which is approximable from above and such that } g(n) \leq f(n) \text{ for all } n \in \mathbb{N}. \tag{5.5}$$

We note that (5.5) holds for all $\Delta_n^0, n \in \mathbb{N}$, but we are only interested in Δ_2^0 here.

Proof of (5.5). Let $f(n)[s]$ be a computable approximation to f . Without loss of generality, we can assume that for all stages s and all $n \leq m \leq s$, we have $f(n)[s] \leq f(m)[s]$. Let $g(n)[s] = \min\{f(n)[t] \mid n \leq t \leq s\}$ for each $n \leq s$. Clearly $g(n) = \lim_s g(n)[s]$ is Δ_2^0 and $g(n) \leq f(n)$ for all $n \in \mathbb{N}$. Also, g is non-decreasing. To show that it is unbounded, let $c, n, s_0 \in \mathbb{N}$ such that $f(n)[s] > c$ for all $s \geq s_0$. Clearly $g(s_0) > c$. \square

Case (a) in Theorem 5.2 is due to Frank Stephan (see [21, Theorem 5.2.25]). As we explain below, Case (b) follows from a combination of Stephan’s theorem and (5.5). For completeness, we give the full argument, the second part being along the lines of the proof of [21, Theorem 5.2.25].

Theorem 5.2. *Suppose that $f : \mathbb{N} \rightarrow \mathbb{N}$ is Δ_2^0 and $\lim_n f(n) = \infty$. If f satisfies one of the following*

- (a) *it can be computably approximated from above*
- (b) *it is non-decreasing*

then there is a Turing complete c.e. set which obeys f . In particular, f is not a gap function for K -triviality of c.e. sets.

Proof. If g satisfies (a) and the assumptions of the theorem, then the non-decreasing function $h(n) = \min\{g(i) \mid i \geq n\}$ also does and $h(n) \leq g(n)$ for all $n \in \mathbb{N}$. So g bounds a function satisfying (a), (b) and the assumptions of the theorem. Moreover by (5.5), any function satisfying (b) and the assumption of the theorem bounds a function with the same properties which also satisfies (a). Hence to prove the theorem it suffices to show that given any Δ_2^0 unbounded non-decreasing function which has an approximation from above, there is a Turing complete c.e. set which obeys it. Let f be such a function with an approximation $f(n)[s]$ such that $f(n)[s] \leq f(n+1)[s]$ and $f(n)[s] \leq f(n)[s+1]$ for all $s \in \mathbb{N}$ and $n \leq s$. In the following, for each real number x , let $\lfloor x \rfloor$ denote the largest integer m such that $m \leq x$. Construct a c.e. set A as follows.

Construction. At stage $2s+1$, find the least $n < s$ such that the number k of zeros in $A[2s] \upharpoonright_n$ is larger than $\lfloor f(n)[s]/2 \rfloor$, and enumerate $n-1$ into A . At stage $2s+2$, if m is the least number $< s$ enumerated in \emptyset' at stage s switch the m th zero position of $A[2s+1]$ into 1.

Verification. By the properties of the approximation $f(n)[s]$, we have that whenever $n-1$ is enumerated into A at stage $2s+1$, for each $i \leq n$ the number of zeros in $A[2s] \upharpoonright_i$ is $\leq \lfloor f(i)[s]/2 \rfloor$. Moreover, in this case the number of zeros in $A[2s+1] \upharpoonright_n$ is exactly $\lfloor f(n)/2 \rfloor$ (otherwise there would not have been such an enumeration). By the construction, if m is the least number enumerated in \emptyset' at stage s and n is enumerated into A at stage $2s+2$, then $A[2s+2] \upharpoonright_n$ has $m-1$ zeros.

First, we show that $\mathbb{N} - A$ is infinite. If A was finite, there is nothing to show. Otherwise there are infinitely many stages t where some number n is enumerated into A and $A[t] \upharpoonright_n = A \upharpoonright_n$. According to the above remarks, for each of these numbers n :

- if n was enumerated at stage $2s+1$, $A \upharpoonright_n$ has $\lfloor f(n)[s]/2 \rfloor$ zeros
- if n was enumerated at stage $2s+2$, then $A \upharpoonright_n$ has $m-1$ zeros, where m is the least number enumerated in \emptyset' at stage s .

Since $f(n)$ is unbounded (and the approximation is non-increasing) it follows that A has infinitely many zeros (i.e. $\mathbb{N} - A$ is infinite). In order to compute $\emptyset'(m)$ from A , we just need to search for n, s such that $A \upharpoonright_n$ has m zeros and $A[s] \upharpoonright_n = A \upharpoonright_n$. Then by the construction $\emptyset'(m) = 1$ iff $\emptyset'(m)[s] = 1$.

Second, we show that A has the required initial segment complexity. By the construction, an induction shows that the number of zeros in $A \upharpoonright_n$ is $\leq \lfloor f(n)/2 \rfloor$ for all $n \in \mathbb{N}$. Therefore, $K(A \upharpoonright_n) \leq^+ K(n) + \lfloor f(n)/2 \rfloor + K(f(n))$ since to describe $A \upharpoonright_n$, we only need to know $n, f(n)$ and a string of length $\lfloor f(n)/2 \rfloor$ indicating the digits in $A \upharpoonright_n$ that are not 1 at the point of the construction where the number of 1s in the string $A \upharpoonright_n$ is $\lfloor f(n)/2 \rfloor$. Since $K(m) <^+ m/2$ for all numbers m , we have $K(A \upharpoonright_n) \leq^+ K(n) + f(n)$ for each $n \in \mathbb{N}$. \square

The following result shows that the conditions in Theorem 5.2 are essential.

Proposition 5.3. *There is a Δ_2^0 function f such that $\lim_n f(n) = \infty$ and f is a gap function for K -triviality of Σ_1^0 sets.*

Proof. Let (W_e) be an effective list of all c.e. sets. We meet the following requirements.

R_e : If W_e obeys f with constant e then $K(W_e \upharpoonright_n) \leq K(n) + 2e$ for almost all n .

Since each c.e. set has infinitely many indices, the satisfaction of these requirements implies that f is a gap function for K -triviality of Σ_1^0 sets. We say that R_e requires attention at stage s if $K(W_e \upharpoonright_s) > K(s) + 2e$. This property is decidable in \emptyset' .

At stage s , find the least $e \leq s$ such that R_e requires attention and is not satisfied. Let $f(s) = e$ and say that R_e is satisfied. If there is no such e , let $f(s) = s$. It is easy to verify that all R_e are met, and from some stage on, they are either satisfied or do not require attention. Moreover, since each R_e 'receives attention' at most once, $\lim_n f(n) = \infty$. \square

The following result contrasts Theorem 5.2.

Theorem 5.4. *There is a gap function for K -triviality of Δ_2^0 sets which is unbounded, non-decreasing and can be \emptyset' -computably approximated from above (in particular, it has Σ_2^0 degree).*

Proof. This is similar to the proof of Theorem 3.1, so we just give a sketch. To make f of Σ_2^0 degree, it suffices to define a \emptyset' -computable approximation to it from above. We meet the following conditions:

N_e : $[\Phi_e^{\emptyset'} \text{ is total and } \Phi_e^{\emptyset'} \not\leq_K \emptyset] \Rightarrow \exists n [K(\Phi_e^{\emptyset'} \upharpoonright_n) \not\leq K(n) + f(n) + e]$.

If we had only one N_e to satisfy, we would define $f(n)[s] = n$ while searching (recursively in \emptyset') for some $m > e$ and a stage t such that

$\Phi_e^{\emptyset'}[t] \upharpoonright_m \downarrow$ and $K(\Phi_e^{\emptyset'} \upharpoonright_m) > K(m) + 2e$.

If and when m, t are found, we let $f(i)[t] = e$ for all $i \in [e, m]$ and continue as before, defining $f(n)[s] = n$ for $n \in (m, s]$ and $s > t$. In this case, m is called a witness for N_e . In the global construction, we make sure that N_e can only modify f on arguments that are larger than the largest witness that any $N_i, i < e$ may have. This ensures that f is approximated monotonically from above. So if $k_e[s]$ is the least number, which is larger than any witness of $N_i, i < e$ at stage s and larger than e , strategy N_e at stage s looks for $m \in (k_e[s], s]$ such that

$\Phi_e^{\emptyset'}[s] \upharpoonright_m \downarrow$ and $K(\Phi_e^{\emptyset'} \upharpoonright_m) > K(m) + 2e$. (5.6)

If it finds such, it sets $f(i)[s] = e$ for all $i \in (k_e, m]$. Note that there is no injury amongst different strategies. Moreover f is unbounded as each N_e acts at most once and never sets the values of f below e . For the same reason each N_e 'receives attention' at some stage, or is trivially satisfied. If it lowers the values of f , it is satisfied by (5.6). \square

Proposition 2.3 shows that if a function f has finite \liminf then there is a bound for the complexity of all sets that obey it (they are computable from $f \oplus \emptyset'$). In particular, the class of sets that obey it is countable. By (5.2), the converse does not hold. However we have the following, which can be seen as a generalization of the fact from [23,19] that the \leq_K -lower cone below any random set is uncountable.

Theorem 5.5. *Suppose that for some $f : \mathbb{N} \rightarrow \mathbb{N}$ we have $\lim_n(f(n) - K(n)) = \infty$. Then there is some $c \in \mathbb{N}$ such that class of sets that obey f with constant c is uncountable. In particular, it contains the paths of a perfect pruned $\Delta_2^0(f)$ tree.*

Proof. We use an oracle argument to construct a perfect pruned $\Delta_2^0(f)$ tree T whose infinite paths obey f with constant c . There is a constant c such that for each string σ and all $n, m \in \mathbb{N}$,

$$K(\sigma 0^{m+n+1}) < K(m+n+1+|\sigma|) + K(\sigma) + c \tag{5.7}$$

$$K(\sigma 0^n 10^m) < K(m+n+1+|\sigma|) + K(\sigma) + K(n) + c. \tag{5.8}$$

Indeed, to describe $\sigma 0^{m+n+1}$, we just need a description of σ (which also lets us compute $|\sigma|$) and a description of $m+n+1+|\sigma|$. Similarly, to describe $\sigma 0^n 10^m$ it is enough to have the above and a description of n (so that we know where to put the 1 which is at the $n+1$ digit after the digits of σ). If we look at T as a map from $2^{<\omega}$ to $2^{<\omega}$ (preserving comparability and incomparability relations), level k of the tree consists of the strings $T(\rho)$ for ρ of length k . The strings of level k will have the same length ℓ_k .

Suppose inductively that level k of the tree has already been defined, and for each string σ on that level the sequence $\sigma 0^i$ obeys f with constant c . Note that letting level 0 of T consisting of the empty string, this assumption holds for level 0 (since $K(i)$ is by definition $K(0^i)$ for each $i \in \mathbb{N}$).

For the definition of level $k+1$, find $n > \ell_k$ such that $f(\ell_k+n+1+m) > K(\sigma) + K(n)$ for all $m \in \mathbb{N}$ and each σ on level k of T . For each such σ define its two successors in level $k+1$ to be $\sigma 0^j$ for $j = 0, 1$. By (5.7) and (5.8), we have that for each string τ of level $k+1$, the sequence $\tau 0^i$ obeys f with constant c .

Since $\lim_n(f(n) - K(n)) = \infty$ and $\lim_n K(n) = \infty$, all levels of T will be defined. By induction, all paths of T obey f . Moreover, only Π_1^0 questions were asked during the definition of T . Hence T is Δ_2^0 in f . \square

It may be interesting to examine if the proof of **Theorem 5.5** can be ‘effectivized’, so that we may obtain a perfect $\Pi_1^0(f)$ class of reals that obey f with constant c under the hypothesis that $\lim_n(f(n) - K(n)) = \infty$.

Since there is a Δ_4^0 unbounded non-decreasing gap function K -triviality (see (5.2)) and by **Theorem 5.2**, no Δ_2^0 function can have this property, it becomes interesting to ask for a Δ_3^0 function with this property. This problem was recently solved in [3]. It was shown that there is a Δ_3^0 unbounded non-decreasing gap function for K -triviality. This argument relies on a result in [7] about the arithmetical complexity of the function that gives the number of K -trivial sets with respect to a given constant. In [3], it is also proved that given any Δ_2^0 unbounded non-decreasing function f there exists a constant c and an uncountable collection of reals X which obey f with constant c . This latter result complements **Theorems 5.2** and **5.5** in this paper.

A well known problem in the LK degrees was to obtain a characterization of the oracles which have uncountable lower cones with respect to \leq_{LK} (see [2,15]). This problem was solved in [5] by showing that this class coincides with the low for Ω oracles (i.e. the ones relative to which the halting probability is Martin-Löf random). The same question can be asked about the K -degrees. Note that by **Propositions 2.2** and **2.3**, the cone below a c.e. set is always countable, while by **Theorem 5.5** there are many sets with uncountable lower cone (including all random sets). Another way to ask the same question is the following.

Problem 5.6. Characterize the functions $f : \mathbb{N} \rightarrow \mathbb{N}$ with the property that the class of sets that obey them (with any constant) is countable.

Finally, we would like to suggest that it may be interesting to study the connection between the functions we discussed in this section and the so-called Solovay functions that were studied in [4,11].

Acknowledgements

We would like to thank Adam Day for reading earlier drafts of this paper. The first author was supported by an *International Young Scientist Fellowship* from the Chinese Academy of Sciences and the *Grand project: Network Algorithms and Digital Information* of the Institute of Software, Chinese Academy of Sciences. This work was part of the second author’s M.Sc. Thesis [22] at the University of Amsterdam.

References

- [1] George Barmpalias, Elementary differences between the degrees of unsolvability and the degrees of compressibility, *Ann. Pure Appl. Logic* 161 (7) (2010) 923–934.
- [2] Barmpalias, George, Relative randomness and cardinality, *Notre Dame J. Form. Log.* 51 (2) (2010).
- [3] Baartse Martijn, George Barmpalias, On the gap between trivial and nontrivial initial segment prefix-free complexity (2010) (submitted for publication).

- [4] Laurent Bienvenu, Rod Downey, Kolmogorov complexity and Solovay functions, in: 26th Annual Symposium on Theoretical Aspects of Computer Science, STACS 2009, Dagstuhl Seminar Proceedings LIPIcs 3, 2009, pp. 147–158.
- [5] George Barmpalias, Andrew E.M. Lewis, Chaitin's halting probability and the compression of strings using oracles, Proc. Roy. Soc. 2011 (in press).
- [6] George Barmpalias, Andrew E.M. Lewis, Mariya Soskova, Randomness, lowness and degrees, J. Symbolic Logic 73 (2) (2008) 559–577.
- [7] George Barmpalias, Tom Sterkenburg, On the number of infinite sequences with trivial initial segment complexity. Preprint, September 2010.
- [8] Barbara F. Csima, Antonio Montalbán, A minimal pair of K -degrees, Proc. Amer. Math. Soc. 134 (5) (2006) 1499–1502 (electronic).
- [9] Rod Downey, Denis Hirshfeldt, Algorithmic Randomness and Complexity, Springer, 2010.
- [10] O. Demuth, A. Kučera, Remarks on 1-genericity, semigenericity, and related concepts, Comm. Math. Univ. Carolinae 28 (1987) 85–94.
- [11] Rupert Hölzl, Thorsten Kräling, Wolfgang Merkle, Time-bounded Kolmogorov complexity and Solovay functions. In, in: Mathematical Foundations of Computer Science, 2009, in: Lecture Notes in Comput. Sci., vol. 5734, Springer, 2009, pp. 392–402.
- [12] Bjørn Kjos-Hanssen, Wolfgang Merkle, Frank Stephan, Kolmogorov complexity and the recursion theorem, in: STACS, 2006, pp. 149–161.
- [13] B. Kjos-Hanssen, J. Miller, R. Solomon, Lowness notions, measure, and domination, J. Lond. Math. Soc. 84 (2011).
- [14] Bjørn Kjos-Hanssen, Wolfgang Merkle, Frank Stephan, Kolmogorov complexity and the recursion theorem, Trans. Amer. Math. Soc. 363 (2011).
- [15] Joseph S. Miller, The K -degrees, low for K degrees, and weakly low for K sets, Notre Dame J. Formal Logic 50 (4) (2010) 381–391.
- [16] Joseph S. Miller, André Nies, Randomness and computability: open questions, Bull. Symbolic Logic 12 (3) (2006) 390–410.
- [17] Wolfgang Merkle, Frank Stephan, On C -degrees, H -degrees and T -degrees, in: Twenty-Second Annual IEEE Conference on Computational Complexity, CCC 2007, San Diego, USA, 12–16 June 2007, IEEE Computer Society, Los Alamitos, CA, USA, 2007, pp. 60–69.
- [18] Joseph S. Miller, Liang Yu, On initial segment complexity and degrees of randomness, Trans. Amer. Math. Soc. 360 (6) (2008) 3193–3210.
- [19] Joseph S. Miller, Liang Yu, Oscillation in the initial segment complexity of random reals, Adv. Math. (2010) (in press).
- [20] André Nies, Lowness properties and randomness, Adv. Math. 197 (1) (2005) 274–305.
- [21] André Nies, Computability and Randomness, Oxford University Press, 2009.
- [22] Charlotte Vlek, Definability in the degrees of randomness. M.Sc. Dissertation, University of Amsterdam, 2010.
- [23] Liang Yu, Decheng Ding, Rod Downey, The Kolmogorov complexity of the random reals, Ann. Pure Appl. Logic 129 (1–3) (2004) 163–180.